



U.S. DEPARTMENT OF
ENERGY

Office of
Science

DOE/SC-ARM-16-014

Deployment of ARM Aerial Facility (AAF) Scanning Mobility Particle Sizer Field Campaign Report

J Wang

March 2016



DISCLAIMER

This report was prepared as an account of work sponsored by the U.S. Government. Neither the United States nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the U.S. Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the U.S. Government or any agency thereof.

Deployment of ARM Aerial Facility (AAF) Scanning Mobility Particle Sizer Field Campaign Report

J Wang, Brookhaven National Laboratory
Principal Investigator

March 2016

Work supported by the U.S. Department of Energy,
Office of Science, Office of Biological and Environmental Research

Executive Summary

Atmospheric aerosols influence global climate by scattering and absorbing sunlight (direct effects) and by changing the microphysical structure, lifetime, and coverage of clouds (indirect effects). While it is widely accepted that aerosol indirect effects act to cool the Earth-atmosphere system by increasing cloud reflectivity and coverage, the magnitudes of the indirect effects are poorly quantified. One key aerosol property for understanding aerosol indirect effects is the ability of aerosol particles to form cloud droplets at atmospheric relevant supersaturations—i.e., cloud condensation nuclei (CCN) activity. For particles consisting of typical atmospheric inorganic compounds, their CCN activity is well understood and can be effectively predicted using Köhler theory based on physicochemical properties of the solute, such as its mass, molar volume, and activity coefficient. However, atmospheric aerosols often consist of hundreds of organic species, which can contribute ~20-90% to the total fine aerosol mass. Depending on their properties, organic species can significantly influence the ability of aerosol particles to act as CCN and form cloud droplets. This project focuses on the CCN activity of secondary organic aerosol compounds formed from key biogenic volatile organic compounds (VOCs) under representative conditions, and the relationship between the hygroscopicity and composition of organic aerosols.

The ARM Aerial Facility (AAF) scanning mobility particles sizer (SMPS) was deployed during a ~ 6 week intensive measurement campaign, taking place June 15-July 31 2015 at the Pacific Northwest National Laboratory (PNNL) Environmental Simulation Chamber. The SMPS was operated with a CCN counter. Aerosol particles were first classified by the differential mobility analyzer inside the SMPS; the classified aerosol will then be simultaneously characterized by a condensation particle counter (part of the SMPS) and the CCN counter.

Acronyms and Abbreviations

AAF	ARM Aerial Facility
ARM	Atmospheric Radiation Measurement Climate Research Facility
ASR	Atmospheric System Research
BVOC	biogenic volatile organic carbon
CCN	cloud condensation nuclei
CCNc	CCN counter
CPC	Condensation Particle Counter
DMA	Differential Mobility Analyzer
PNNL	Pacific Northwest National Laboratory
POA	primary organic aerosol
SMPS	Scanning Mobility Particle Sizer
SOA	secondary organic aerosol
VOC	volatile organic carbon

Contents

Executive Summary	ii
Acronyms and Abbreviations	iii
1.0 Background.....	1
2.0 Notable Events or Highlights	1
3.0 Results	1
4.0 Deployment of AAF Scanning Mobility Particle Sizer Publications	2
4.1 Journal Articles/Manuscripts.....	2
4.2 Meeting Abstracts/Presentations/Posters	2

1.0 Background

Atmospheric aerosols influence global climate by scattering and absorbing sunlight (direct effects) and by changing the microphysical structure, lifetime, and coverage of clouds (indirect effects). While it is widely accepted that aerosol indirect effects act to cool the Earth-atmosphere system by increasing cloud reflectivity and coverage, the magnitudes of the indirect effects are poorly quantified. One key aerosol property for understanding aerosol indirect effects is the ability of aerosol particles to form cloud droplets at atmospheric relevant supersaturations—i.e., cloud concentration nuclei (CCN) activity. For particles consisting of typical atmospheric inorganic compounds, their CCN activity is well understood and can be effectively predicted using Köhler theory based on physicochemical properties of the solute, such as its mass, molar volume, and activity coefficient. However, atmospheric aerosols often consist of hundreds of organic species, which can contribute ~20-90% to the total fine aerosol mass. Depending on their properties, organic species can significantly influence the ability of aerosol particles to act as CCN and form cloud droplets. This project focuses on the CCN activity of secondary organic aerosol compounds formed from key biogenic volatile organic compounds (VOCs) under representative conditions, and the relationship between the hygroscopicity and composition of organic aerosols.

The ARM Aerial Facility (AAF) scanning mobility particle sizer (SMPS) was deployed during a 6-week intensive measurement campaign, taking place June 15-July 31 2015 at the Pacific Northwest National Laboratory (PNNL) Environmental Simulation Chamber. The SMPS was operated with a CCN counter. Aerosol particles were first classified by the differential mobility analyzer inside the SMPS; the classified aerosol will then be simultaneously characterized by a condensation particle counter (part of the SMPS) and the CCN counter.

The principle investigator of this deployment is Jian Wang from Brookhaven National Laboratory. Co-investigators include Fan Mei from PNNL and Ryan Thalman from Brookhaven National Laboratory (current affiliation: Snow College).

2.0 Notable Events or Highlights

The deployment of AAF went smoothly, with no unusual observations.

3.0 Results

The deployments of AAF SMPS provided aerosol-size spectrum from 40 to 250 nm, a size range in which CCN activity is strongly influenced by particle composition. The aerosol size spectrum data are now available from the ARM Data Archive.

4.0 Deployment of AAF Scanning Mobility Particle Sizer Publications

4.1 Journal Articles/Manuscripts

Data analysis is still ongoing. A manuscript describing the results from this deployment is planned.

4.2 Meeting Abstracts/Presentations/Posters

“Hygroscopicity of organic aerosols and aerosol indirect effects on climate”, invited platform presentation, 250th American Chemical Society National Meeting, Boston, MA, August, 2015.



U.S. DEPARTMENT OF
ENERGY

Office of Science